



PHYSIOTHERAPY
CLINICAL
GUIDELINES

Huntington's disease

European Huntington's Disease
Network Physiotherapy
Working Group



Clinical Implementation Strategy

People with Huntington's disease (HD) demonstrate a range of physical, cognitive, psychological and social care needs over an extended time frame. One of the difficulties in developing clinical guidelines for complex neurodegenerative diseases such as HD is the heterogeneity of clinical signs and symptoms. Whilst staging of the disease process (e.g. early, middle, late) can provide a general framework for intervention, within each stage there is a wide range of potential impairments that can impact an individual's level of functional activity and life participation. This makes structuring of consistent therapeutic approaches problematic. This problem is not unique to HD, and has been documented in other physiotherapy patient groups, most notably low back and neck dysfunction, and for general neurorehabilitation patient groups [1-3].

Clinical guidelines are evidenced-based recommendations for clinical practice in specific conditions [4, 5]. The availability of clinical guidelines facilitates uniformity of care and standards of practice with the aim of improving quality of care provision. It is critically important for health professionals to be able to define and document their assessment and intervention strategies. The application of appropriate guidelines in practice, as well as systematic outcome evaluation, has the potential to promote evidenced-based delivery of care for the benefit of the person with HD.

Members of the Physiotherapy Working Group (PWG) are advocating a treatment-based classification approach to specifically guide intervention strategies in HD. Utilisation of treatment-based classifications may help to address the heterogeneity of impairments and activity limitations seen in people with HD, and provide a structure for standardised data collection of interventions and outcomes. It is hoped that by creating sub classifications of patients' impairments and problems, and matching those with more specific interventions, therapists may improve outcomes in their patients. The following seven classifications have been developed to better categorize patients with HD, who can present with a wide array of functional problems and physical, cognitive and behavioural impairments. It is the aim of the PWG that these treatment-based classifications will subsequently inform evaluation of complex interventions and advance research into care and evidenced-based service delivery for people with HD. As the complex nature of HD makes it unlikely that any one professional will have all the skills needed for best practice, these classifications have also incorporated referral to relevant members of the multi-disciplinary team

Therapists should review the signs and symptoms of each classification to determine the best fit for a particular patient. Once a classification has been determined, therapists should use it as a guide to selecting appropriate evaluation measures and intervention strategies. Importantly, these classifications are not designed to take the place of independent clinical decision-making based on each patient's particular signs and symptoms, but rather should provide a framework for more consistent patient management through the disease spectrum.

Classification	Description	Stages
A. Exercise Capacity and Performance	Absence of motor impairment or specific limitations in functional activities; potential for cognitive and/or behavioural issue	Pre-manifest/early
B. Planning and sequencing of tasks (including bradykinesia)	Presence of apraxia or impaired motor planning; slowness of movement and/or altered force generation capacity resulting in difficulty and slowness in performing functional activities.	Early-mid
C. Mobility, Balance and Falls Risk	Ambulatory for community and/or household distances; impairments in balance, strength or fatigue resulting in mobility limitations and increased falls risk	Early-mid
D. Secondary adaptive changes and deconditioning	Musculoskeletal and/or respiratory changes resulting in physical deconditioning, and subsequent decreased participation in daily living activities, or social work environments.	Early-mid
E. Impaired postural control and alignment in sitting	Improper alignment due to adaptive changes, involuntary movement, muscle weakness and incoordination resulting in limitations in functional activities in sitting.	Mid-late
F. Respiratory dysfunction	Impaired respiratory function and capacity; limited endurance; impaired airway clearance resulting in restrictions in functional activities and risk for infection	Mid-late
G. End Stage Care	Active and passive range of motion limitations and poor active movement control resulting in inability to ambulate; dependent for most ADLs; difficulty maintaining upright sitting position	Late

A. EXERCISE CAPACITY AND PERFORMANCE

Description : Absence of motor impairment or specific limitations in functional activities; potential for cognitive and/or behavioural issues / **Stage :** Pre-manifest/early

Signs and symptoms / key issues and potential issues	General aims	Treatment options
<p>Participation : possible quality of life changes</p> <p>Activities : no problems</p> <p>Impairments : Potential for: early gait changes, poor endurance or fitness; mild chorea; cognitive and/or behavioural issues; poor endurance and limited physical activity ; lack of motivation and/or apathy ; anxiety and/or depression [6, 7] ; sleep disturbance which may exacerbate the above impairments</p>	<p>Facilitate health education and advice - this includes general health promotion strategies, referral for exercise on a prescriptive plan, and agenda setting to optimise exercise performance.</p> <p>Patient & family education – focus on importance of early intervention in HD. In animal models of HD, early enrichment of the environment (i.e., exercise) was shown to delay symptom onset and severity [8]. Increased physical activity in HD mice was critical for successful outcomes of disease modifying treatments such as cell transplants and reconnection of grafted tissue. [9].</p> <p>Facilitate active lifestyle – this is thought to delay the disease onset in people with HD [10] and to reduce the risk of other neurodegenerative disease such as Alzheimer's disease [11] and Parkinson's disease [12]. In addition, participation in regular exercise has potential to result in improved strength, cardiovascular fitness, exercise tolerance, functionality, mobility and mood.</p>	<p>Baseline testing for fitness level should be completed prior to exercise prescription. Consider education on fatigue and the timing of intervention/exercises during the day as well as careful instruction on safety during exercise.</p> <p>Identify barriers and facilitators to initiate and maintain an exercise programme [13]. Techniques such as behavioural motivation can be used to help patient identify barriers and facilitators and explore strategies to manage them [14]. Involvement of a caregiver/friend/spouse can make the programme more successful.</p> <p>Gym based exercises can be of benefit to physically-able individuals, as well as people with long term neurologic conditions such as Parkinson's Disease [15]. Small case reports in HD [16, 17] indicate that if properly supported, people with HD can enjoy the health benefits of physical activity. Interestingly, there is underutilization of PT services in early-stage HD [18].</p> <p>Exercise in early stage HD should include:</p> <ul style="list-style-type: none"> • Individualized goal setting and home exercise programme prescription for optimization of services in a life-long disease process. • A focus on task-specific functional activities incorporated into the exercise programme. • A warm-up and cool-down. • Careful monitoring of vital signs, dyspnoea, fatigue, pallor, dizziness and specific HD-related signs at rest, during and after exercise <p>Frequency, intensity, duration and mode are dependent on the baseline fitness level of the individual; however, focus should be on the ACSM goal of exercise for both aerobic and strength training. [19].</p> <p>Aerobic exercise : Frequency : 3 to 5 times a week; Intensity: 65% to 85% of the maximal heart rate; 55% to 65% of maximal heart rate for de-conditioned individuals ; Duration : at least 30 minutes of continuous or intermittent training per day (minimum of 10-minute bouts accumulated throughout the day); Mode : any activity that the individual enjoys that uses large muscle groups which can be maintained continuously and is rhythmic and aerobic in nature (e.g. walking, jogging, swimming, and biking).</p> <p>Resistance exercises : Frequency & Duration : 8 to 12 repetitions per exercise; One set of 8 to 10 exercises that conditions large muscle groups 2 to 3 times a week ; Intensity : 65% to 70% of 1 rep max for upper body and 75% to 80% of 1 rep max for lower body parts ; Mode : Resistance training should be progressive and individualized.</p> <p>Exercise programme ideas : walking (treadmill and over-ground), stationary bicycling, horse-back riding, strength-training, balance training, core stability training, video-game based exercise (Nintendo Wii, Dance Dance Revolution) [20]. Yoga, pilates, tai chi and relaxation are also recommended.</p>

Outcome Measures :

PAR-Q : The Physical Activity Readiness Questionnaire (PAR-Q) can be utilised as an initial screening method to determine any contraindications to exercise. If any reason to doubt safety in participating in exercise, the physiotherapist should refer persons to their general practitioners (GPs) for a full assessment [21].

IPAQ : International Physical Activity Questionnaire: is a useful questionnaire for obtaining information about weekly activity levels at home, work and leisure. It is reliable and valid in 18-65 year old healthy adults as an epidemiological measure. It is not valid as a measure of change or to assess the effects of an intervention [22].

Activity Monitor : if available, a useful measure of physical decline in HD as hypokinesia is related to functional capacity in HD. May also be used to show change in activity over time [23-26].

Borg RPE (or CR-10) : is a useful objective measure of perceived exertion. In addition, it allows patients to self-monitor during their exercise routine. This RPE (rate of perceived exertion) scale has become a standard method for evaluating perceived exertion in exercise testing, training and rehabilitation, and has been validated against objective markers of exercise intensity [27, 28].

Goal Attainment Scale : assists with appropriate goal setting, quantifying achievement, and even weighting of goals to reflect the personal importance of the goal to the patient and/or the opinion of the therapist on the difficulty of achieving the goal [29].

6 minute walk test (6MWT) for measuring endurance and cardiovascular fitness [30]

Interdisciplinary :

Nutrition: assessment and management of caloric intake, dietary needs and changes to diet when exercising.

Psychology: assessment and management of behavioural issues.

Occupational Therapy: assessment and management of ADL-related deficits.

Speech & Language Therapy: for clear communication of printed exercise routines.

B. PLANNING AND SEQUENCING OF TASKS

Description : Presence of apraxia or impaired motor planning; slowness of movement and/or altered force generation capacity resulting in difficulty and slowness in performing functional activities. / **Stage :** Early-mid

Signs and symptoms / key issues and potential issues	General aims	Treatment options
<p>Participation : Patients may note a decline in independence in their daily routine, and may experience difficulty managing previously automatic tasks such as rising from a chair or walking.</p> <p>Activities : Difficulty and slowness in performing functional activities (dressing, bathing, stair climbing, indoor/outdoor ambulation, ADLs)</p> <p>Impairments :</p> <ul style="list-style-type: none"> • Apraxia – impaired motor planning [31] • Movement speed – slowed speed of movement during purposeful task • Force generation – generalized and/or specific muscle weakness ; delayed onset of muscle response • Safety awareness and insight into deficits 	<p>Creation of individualized patient-centred goals that focus on the specific impairments of the patient.</p> <p>Examples include :</p> <ul style="list-style-type: none"> • Improve ability to perform functional tasks • Increase speed of movement • Maximize safety 	<p>Task-specific training to address planning and sequencing deficits.</p> <p>Could include :</p> <ul style="list-style-type: none"> • strategy training in daily living activities : this technique teaches internal (for example, the patient is taught to verbalise and implement the task steps at the same time) or external (for example, when aids are used to overcome a functional barrier) compensatory strategies that enable a functional task to be completed [32]; spaced retrieval and errorless learning techniques can guide training of motor activities and learning of skilful activities [33]. • sensory stimulation : deep pressure and soft touch are applied to the patients' limbs; therapies utilizing a multi-sensory stimulation approach noted significant improvements in mood and stimulation which were cumulative over sessions when compared to the control group [34]. • cueing : visual, verbal or physical prompts: enable task completion. Verbal prompts can include attentional strategies with external cues and/or, attentional strategies with internal cues • chaining (forward or backward) : the task is broken down into its component parts. Using backward chaining the task is completed with facilitation from the therapist apart from the final component, which the patient carries out unaided. If successful, further steps are introduced in subsequent trials.

Outcome Measures :

SF-36 : A quickly and easily administered Quality of Life measure commonly used in HD; robust construct validity and test-retest reliability have been demonstrated [35]

10 Meter Walk : used as a measure of gait speed [36].

Physical Performance Test (PPT) : useful for measuring ability to perform everyday tasks including utensil use, writing, donning/doffing a jacket, walking and stair climbing.

Goal Attainment Scale : see A: Exercise Capacity and Performance for description.

Timed Up & Go : [37] The participant is instructed to stand from a chair, walk 10 feet, return to the chair and be seated. Times greater than 13.5 seconds accurately predicted elderly fallers with 90% accuracy. The TUG has been validated in HD. [26, 37]

- TUG Manual complete TUG while carrying a full cup of water; a time difference of >4.5 seconds between TUG and TUG Manual indicated increased risk of future falls in elderly
- TUG Cognitive [38] (Gait speed; dual-task ability) complete the TUG while counting backwards from a randomly selected number between 20 and 100.

Four Square Step Test (4SST) : participants step forward, sideways and backward over low obstacles (usually canes) in a sequence; a useful measure of balance, stepping ability and motor planning [39].

The Apraxia Test : consists of two subtests assessing the ability to use objects or pantomime use and the ability to imitate gestures [40]

Timed Sit-to-Stand Test : patients are asked to stand up and sit back down 5 times from a chair that reaches the head of their fibula. Time to complete 5 repetitions are recorded. The Sit-to-Stand test is commonly used to assess leg strength and balance and has been shown to be a reliable and valid measure in older adults and other patient populations [41] .

Interdisciplinary :

Occupational Therapy: problem solving and strategies for successful completion of activities of daily living. Compensatory strategies for managing deficits in ocular movements [42] and early cognitive impairments.

Speech and Language Therapy: assessment and management of communication (dysarthria and apraxia of speech; influence of cognitive ability in particular difficulties in executive functioning and in working memory on linguistics, speech and communication skills); assessment and management of swallowing and dysphagia and any association with motor function/ planning problems.



C. MOBILITY, BALANCE AND FALLS RISK

Description : Ambulatory for community and/or household distances; impairments in balance, strength or fatigue resulting in mobility limitations and increased falls risk / **Stage :** Early-mid

Signs and symptoms / key issues and potential issues	General aims	Treatment options
<p>Participation :</p> <ul style="list-style-type: none"> • Fear of falling may result in more unwillingness to participate in home, work, and community activities • Difficulty in participating in recreational sports (e.g., cycling, running, soccer, basketball) that require balance and mobility <p>Activities :</p> <ul style="list-style-type: none"> • Difficulty walking in certain environments (i.e., open environments) • Difficulty walking backwards or side-wards • Difficulty turning and changing directions • Difficulty getting in and out of chairs and beds due to vaulting, poor eccentric control • Difficulty with walking while doing a secondary cognitive or motor task due to attentional deficits [43] • High falls risk [26, 44] <p>Impairments :</p> <ul style="list-style-type: none"> • Bradykinesia [45] • Dystonia – affecting trunk (lateral shift; extension), ankles/feet (inversion) [46] • Chorea*/ rigidity [47-49] • Muscle weakness or impaired force production [50] • Impaired motor control (i.e., force modulation deficits causing sudden exaggerated movement changes; impaired eccentric motor control) • Decreased limb coordination resulting in step asymmetry • Fatigue and its associated influence on physical performance may lead to falls • Balance deficits (increased sway in stance and during functional tasks of daily living, delayed responses to perturbations, difficulty with tandem standing and walking) [16, 51, 52] • Gait impairments (decreased speed; stride length; stride width, increased variability in gait parameters)[53, 54] • Cognitive and behavioural issues including not recognizing their own disabilities and doing unsafe behaviours due to impulsiveness, attentional deficits and problems with dual tasking [44, 55, 56] • Deficits in spatial perception causing people with HD to run into walls or tables [57] • Visual disturbances such as difficulty with saccadic eye movements and smooth pursuit can impact balance and walking [58] 	<ul style="list-style-type: none"> • Improve mobility status (increase independence; increase speed; increase distance walked) • Reduce risk of falls or actual falls** • Maintain independent mobility including transfers and walking for as long as possible • Reduce fear of falling which in itself may cause inactivity <p>Outcome Measures : SF 36 : see B Function Assessment section of the UHDRS : which consists of the Functional Capacity Scale, the Independence Scale, and a checklist of common daily tasks for assessment of basic ADLs and IADLs. The total score on the Functional Capacity Scale is reported as the total functional capacity (TFC) score. The Independence Scale is rated from 0 to 100. Higher scores indicate better functioning [65] HD-ADL : an informant rated instrument designed to follow disease progress. An ADL total score is calculated by summing all of the values for the five domains of Personal Care, Home Care, Work and Money, Social Relationships, and Communication. Validity and reliability has been demonstrated for subjects with HD on the seventeen items of adaptive functioning. Scores range from 0 (independent) to 24 (maximum disability) [66] 10m walk : see B TUG : see B 6MWT : see A ABC scale : measures of confidence performing various daily skills; participant rating scale [67] Berg Balance Scale : Measures balance on 14 tasks; cutoff for falls risk in HD is a score of 40 [68, 69] Tinetti Mobility and Gait Test : Measures basic balance and walking ability [70,71] Gait spatial and temporal measures : using GaitRite [72] Falls history UHDRS motor section [65] Assessments of different devices and equipment used during functional activities including the safety during use, alignment, fit, and the patient's ability to care for the devices or equipment Interdisciplinary : Interdisciplinary : nursing, OT, neuropsychologist, neurologist *see Management of chorea **See Policy on Falls and Mobility, EHDN Physiotherapy Working Group</p>	<ul style="list-style-type: none"> • Impairment exercises : strengthening; general conditioning; endurance; range of motion activity to counteract effects of dystonia; coordination exercises; teach strategies to help people with HD identify when fatigue would increase their risk of falls [58, 59] • Balance training to practice the maintenance of postural control in a variety of tasks and environments <ul style="list-style-type: none"> - Train patients to step in response to perturbations in all directions with speed and accuracy - Practice activities that require automatic responses (e.g., throwing ball) to elicit postural responses and train faster movements - Progress activities from wide to narrow BOS , static to dynamic activities, low to high COG , increasing degrees of freedom - Task-specific practice of functional activities such as transfers, reaching high and low, stair climbing, etc. to train balance control during activities of daily living • Task specific training to address walking tasks, ideally in specific environments (e.g. outdoor; obstacles) ; external cueing [16, 59, 60] <ul style="list-style-type: none"> - Train patients to walk forwards, backwards, and sideways at different speeds and over different surfaces - Use metronome [55, 61, 62], lines on floor to promote step initiation, bigger steps, faster speed, and gait symmetry - Teach strategies as to how to get up from floor if they fall - Teach safety awareness and adaptation of environment (reduce clutter, slippery surfaces, loose rugs, poor lighting, sharp or breakable objects), furniture - Provision of assistive devices (4 wheeled walker with brakes) when appropriate; if patient is unsafe with assistive devices, human support such as holding the person's arm may be helpful [63] <ul style="list-style-type: none"> - Family/carer education for guarding and/or assistance during ambulation • Adaptive devices/equipment <ul style="list-style-type: none"> - Wheelchair prescription (long distance mobility) - Shoe/orthotic evaluation (shoes with ankle support such as high top tennis shoes or boots; heel wedge and/or lateral wedge for ankle dystonia in inversion/eversion direction; ankle foot orthosis for ankle dystonia in dorsiflexion/plantarflexion direction; custom made shoe inlay for individuals with clawing of toes during walking) - Prescribe protective gear – helmets, elbow/knee pads to be worn by person with HD at risk for falls • Teach compensatory strategies for cognitive impairments and inability to dual task <ul style="list-style-type: none"> - Teach patients to focus their attention on maintaining balance before doing a task that challenges their balance [64] - Have people with HD practice two activities at same time under various practice and context conditions in early stages [55, 56] - Teach patients to break down complex tasks into simpler tasks and attend to one task at a time in middle to late stages

D. SECONDARY AND ADAPTIVE CHANGES AND DECONDITIONING

Description : Musculoskeletal and/or respiratory changes resulting in physical deconditioning, and subsequent decreased participation in daily living activities, or social work environments. / **Stage :** Early/mid

Signs and symptoms / key issues and potential issues	General aims	Treatment options
<p>Participation :</p> <ul style="list-style-type: none"> • Decreased level of physical fitness • Decreased participation in ADLs, social or work environments <p>Activities :</p> <ul style="list-style-type: none"> • Decreased daily walking and physical activity levels in people with HD compared to healthy individuals [24] • Daily walking levels significantly more reduced in people with HD who were recurrent fallers compared to those who were non-fallers [26]. <p>Impairments :</p> <ul style="list-style-type: none"> • Musculoskeletal changes - loss of ROM, loss of strength due to inactivity [50] • Respiratory changes – reduced endurance • Cognitive problems such as memory deficits, loss of initiative or insight into problems [43] • Psychological issues such as depression, apathy, anxiety • Weight loss due to multiple factors may contribute to weakness, fatigue [73] • Pain caused by dystonia, muscle imbalances, trauma from falls or hitting objects, immobility [74] • Balance and gait impairments resulting in frequent falls (see Mobility, Balance, and Falls Risk) 	<ul style="list-style-type: none"> • Prevent further physical, cognitive, and psychological deterioration • Improve strength, balance strategies, and stamina • Motivate patients to regain some control of their lives by adopting a healthy lifestyle • Increase patient and caregiver awareness of benefits of regular exercise and detrimental effects of inactivity • Weight control • Pain management 	<ul style="list-style-type: none"> • Educate patient and caregivers on benefits of exercise for people with HD and negative consequences of inactivity • Initiate maintenance programme to prevent secondary adaptive changes and deconditioning [75, 76] • Encourage person with HD to start/restart exercise programme and provide exercise log or diary to record progress • Consider formal or informal programmes : individualized or group exercise programmes in community/hospital settings [60]; exercise video or written instructions with pictures to increase home exercise compliance • Assess for appropriate walking aids, devices (e.g. assistive, adaptive, orthotic) • Treat balance and gait impairments, fear of falling that may underlie activity limitations [77] • Educate caregivers on strategies to motivate (incorporate exercise into daily routine, positive reinforcement, participate in exercise with the patient, select physical activities that the patient enjoys) and assist their loved ones (i.e., cueing, guarding, use of gait belt) with exercise programmes [78] • Encourage patients to seek out enriched environments that are physically and cognitively stimulating and promote social interactions • Teach breathing exercises to maintain full respiratory function • Educate patient on importance of proper nutrition and maintenance of adequate weight • Manage pain appropriately with modalities [79], ROM exercises, proper positioning, protection from injuries, medications, etc.  

Outcome Measures :

- SF-36 (see A)
- Berg Balance Scale (see C)
- 6MWT (see A)

Impairment measures : Aerobic capacity during functional activities, or during standardized exercise test (early stages); Cardiovascular and pulmonary signs and symptoms in response to exercise or increased activity; Weight measurements; Mini Mental State Examination (MMSE)[80]; manual muscle testing or hand held dynamometry for muscle strength; Pain numerical rating scale, Pain Visual Analog scale (VAS), Wong-Baker FACES pain rating scale [74]; UHDRS Behavioral assessment section [65]; Goniometry, end feel assessment, and multisegment flexibility tests for ROM assessment; Respiratory rate, rhythm, and pattern, auscultation of breath sounds, cough effectiveness testing, vital capacity (VC) testing in supine and upright positions or forced vital capacity (FVC) testing for respiratory assessment

Interdisciplinary :

nutritionist, OT, psychologist, personal trainers

E. IMPAIRED POSTURAL CONTROL AND ALIGNMENT IN SITTING

Description : Improper alignment due to adaptive changes, involuntary movement, muscle weakness and incoordination resulting in limitations in functional activities in sitting / **Stage :** mid/late

Signs and symptoms / key issues and potential issues	General aims	Treatment options
<p>Participation :</p> <ul style="list-style-type: none"> • Increased caregiver burden • Withdrawal from society <p>Activities :</p> <ul style="list-style-type: none"> • Difficulty with ADLs including washing, dressing • Difficulty with feeding and swallowing • Inability to stand or sit independently <p>Impairments :</p> <ul style="list-style-type: none"> • Inappropriate musculoskeletal alignment • Decreased range of movement (active and passive) • Soft tissue adaptive changes • Altered base of support leading to:- • Changes in dystonia/ chorea • Poor balance • In-coordinated movement • Pressure areas due to shearing • Potential for falls and soft tissue damage • Risk of aspiration and respiratory complications 	<ul style="list-style-type: none"> • Prevent or limit soft tissue adaptive changes • Minimize risk of infection and skin breakdown • Maximize independence in sitting and facilitate appropriate positioning • Maximize functional ability – eating, reaching tasks, dressing, self positioning, transfers 	<ul style="list-style-type: none"> • Manual handling and falls risk assessment review • Institute positioning schedule for patients with limited active movement • Musculoskeletal realignment and facilitation of more efficient patterns of movement [81]. <p>Options include:</p> <ul style="list-style-type: none"> • Stretches where appropriate • Active and passive range of motion exercises • Positioning (24 hours consideration) • Splinting • Strengthening • Specific trunk stability exercises <ul style="list-style-type: none"> • Assess suitability of seating ; wheelchair evaluation • Respiratory assessment as appropriate [82] • Educate carers and patient re : risk of aspiration • Alignment in each posture [81]

Outcome Measures :

Carer Burden Scale: Used to assess burden of care among caregivers. It is a 29-item scale designed to measure feelings of burden experienced by caregivers of elderly persons with senile dementia [83].

Goal Attainment Scale (see A): decreased dependence in ADL, time tolerated in chair

Interdisciplinary :

Nursing, SALT for swallow assessment, Respiratory physiotherapist, OT, Liaise with specialist OT re: seating and hoist equipment



F. RESPIRATORY DYSFUNCTION

Description : Impaired respiratory function and capacity; limited endurance; impaired airway clearance resulting in restrictions in functional activities and risk for infection / **Stage :** mid/late

Signs and symptoms / key issues and potential issues

Participation :

Participation in social activities e.g. shopping, family activities
Participation in exercise activities

Activities :

Decreased exercise tolerance; limited ability to perform ADLs, ambulation

Impairments :

- breathlessness on exertion or at rest
- decreased exercise capacity
- ineffective cough
- retained secretions
- increased work of breathing
- cyanosis
- decreased oxygen saturation
- dystonia of trunk muscles

General aims

- optimize respiratory function for functional activities
- optimise cardiorespiratory function
- maintain PCF ≥ 270 L/min when well ; PCF ≥ 160 L/min when unwell with cold/respiratory infection [84]
- optimize secretion clearance

Treatment options

- functional exercise
- positioning to manage breathlessness [85]
- breathing exercises; maximal insufflation/exsufflation; glossopharyngeal breathing[84, 86]
- airway clearance techniques [87]
- postural management [88]
- relaxation
- consultation for appropriate walking aids e.g. wheeled walker/rollator
- cardiovascular exercise training [86]

Outcome Measures :

Borg breathlessness/MRC scale breathlessness scale (Australian Lung Foundation)
6MWT: (see A)
Peak cough flow (PCF) [84]
Auscultation/observation/saturation monitor
Forced Vital Capacity

Interdisciplinary :

SALT [89], nursing, respiratory therapist, pulmonologist



G. END STAGE CARE

Description : active and passive range of motion limitations and poor active movement control resulting in inability to ambulate; dependent for most ADLs ; difficulty maintaining upright sitting position / **Stage :** Late

Signs and symptoms / key issues and potential issues	General aims	Treatment options
<p>Participation : complete dependence in functional skills; social isolation</p> <p>Activities : Unable to ambulate; dependent for most ADLs ; difficulty maintaining upright sitting position</p> <p>Impairments :</p> <ul style="list-style-type: none"> • limited volitional control of limbs and trunk • chorea and/or rigidity • limitations in passive range of motion • risk for aspiration/respiratory infection • risk for pressure sores; pain due to positioning or pressure sores or contractures • difficulties or inability to communicate • depressed mood or depression • pain 	<ul style="list-style-type: none"> • Minimize risk of aspiration/respiratory infection • Minimize risk of bed sores • Promote ability to maintain upright sitting position • Promote optimal comfort in bed • Maintain/increase range of motion • Maintain existing ADLs • Support and brief nursing staff or caregivers • If necessary: provide support to organize external help (home-nursing services) or relocation to a specialized institution 	<p>Positioning [75]</p> <ul style="list-style-type: none"> • In bed – utilize pressure relieving mattresses and cushions for optimal positioning; utilize positioning schedule in bed to promote position change (side-lying and supine) • Upright – consultation for appropriate supportive chair. Key features include padded supports (to prevent injury secondary to involuntary movements), tilt in space to maintain appropriate hip angle and allow for change in position for pressure relief; adequate trunk and head supports; adequate padded foot supports • Close cooperation with nursing staff / caregiver to improve transfers, eating position, communication, washing and dressing and to preserve whatever independence the patient has in terms of ADL's. Advise as to signs of aspiration. <p>Range of motion [75]- create range of motion exercise plan to be performed daily by nursing staff, aides, or family members. Regular physiotherapy sessions advisable.</p> <p>Active movement [75] – if able, encourage upright standing with support (consider use of standing table); sitting on edge of bed (with support); active exercises in bed to prevent muscle wasting and prevent breakdown; work with existing capabilities to maintain current ADLs</p> <p>Respiratory (see Respiratory TC)</p>

Outcome Measures :

Braden Risk Assessment Scale: grading tool for risk of pressure ulcers [90]
National Pressure Ulcer Advisory Panel (NPUAP) - pressure ulcer staging [91]; respiratory function assessment; ability to sit upright in adaptive chair for measured period of time; LE/UE range of motion assessment using goniometer; pain assessment with FACES pain scale [74]; Caregiver Burden Scale [83] if patient remains at home

Interdisciplinary :

Nursing, respiratory therapy, SALT, clergy for spiritual support, psychologists for grief counselling, OT for seating, social workers for living wills, power of attorney



References

- Fritz, JM, Brennan GP: Preliminary examination of a proposed treatment-based classification system for patients receiving physical therapy interventions for neck pain. *Physical Therapy* 87(5),513-24 (2007).
- Scheets PL, Sahrman SA, Norton BJ: Use of movement system diagnoses in the management of patients with neuromuscular conditions: a multiple-patient case report. *Physical Therapy* 87(6),654-69, (2007).
- Bello-Haas, VD: A framework for rehabilitation of neurodegenerative diseases: planning care and maximizing quality of life. *Neurology Report* 26(3), 115-129 (2002).
- Ramaswamy B, Ashburn A, Durrant K et al: Quick reference Cards (UK) and guidance notes for physiotherapists working with people with PD. *Parkinson Disease Society* (2009).
- Royal Dutch Society for Physical Therapy: KGNF Guidelines for Physical Therapy in Patients with Parkinson's Disease. *Dutch Journal of Physiotherapy* 114, Supplement 3 (2004).
- Vaccarino AL, Silts T, Anderson KE et al: Assessment of Day-to-Day Functioning in Prodromal and Early Huntington Disease. *PLoS Curr*, 3, RRN1262 (2011).
- Morton AJ, Wood NI, Hastings MH, Hurelbrink C, Barker RA, Maywood ES: Disintegration of the sleep-wake cycle and circadian timing in Huntington's disease. *J Neurosci* 25(1), 157-63 (2005).
- van Dellen A, Cordery PM, Spiers TL, Blakemore C, Hannan AJ: Wheel running from a juvenile age delays onset of specific motor deficits but does not alter protein aggregate density in a mouse model of Huntington's disease. *BMC Neurosci*, 9,34 (2008).
- Dobrossy MD, Dunnett SB: Training specificity, graft development and graft-mediated functional recovery in a rodent model of Huntington's disease. *Neuroscience* 132(3), 543-552 (2005).
- Trembath MK, Horton ZA, Tippett L et al: A retrospective study of the impact of lifestyle on age at onset of Huntington disease. *Mov Disord*. 25(10), 1444-50 (2010).
- Scarmeas, N, Levy G, Tang MX, Manly J, Stern Y: Influence of leisure activity on the incidence of Alzheimer's disease. *Neurology* 57(12), 2236-42 (2001).
- Thacker, EL, Chen H, Patel AV et al: Recreational physical activity and risk of Parkinson's disease. *Mov Disord* 23(1), 69-74 (2008).
- Quinn L, Busse M, Khalil H, Richardson S, Rosser A, Morris H et al: Client and therapist views on exercise programmes for early-mid stage Parkinson's disease and Huntington's disease. *Disability and Rehabilitation* 32(11), 917-28 (2010).
- van Nimwegen M, Speelman AD, Smulders K et al: Design and baseline characteristics of the ParkFit study, a randomized controlled trial evaluating the effectiveness of a multifaceted behavioral program to increase physical activity in Parkinson patients. *BMC Neurol* 10, 70 (2010).
- Goodwin V, Richards SH, Taylor RS, Taylor AH, Campbell JL: The effectiveness of exercise interventions for people with Parkinson's disease: A systematic review and meta-analysis. *Movement Disorders* 23(5), 631-40 (2008).
- Quinn L, Rao A: Physical therapy for people with Huntington disease: current perspectives and case report. *Neurology Report* 26(3), 145-53 (2002).
- Meaney, A, Busse M, Dawes H, Rosser A: Response to a structured exercise programme for Huntington's Disease; a single case study. *British Association of Sports and Exercise Medicine Journal of Sports Science* (2008).
- Busse M, Khalil H, Quinn L, Rosser A: Physical Therapy Intervention for People With Huntington Disease. *Phys Ther* 88(7), 820-31 (2008).
- American College of Sports Medicine. Guidelines for exercise testing and prescription. 8th edition, Philadelphia: Lippincott Williams & Wilkins (2009).
- Kloos A, Kostyk S, Kegelmeyer D: The effect of video game-based exercise on dynamic balance, mobility and UHDRS neuropsychiatric test scores in individuals with Huntington's Disease. *The Journal of the American society for experimental neurotherapeutics* 8(1), 139 (2011).
- Thomas S, Reading J, Shephard RJ: Revision of the Physical Activity Readiness Questionnaire (PAR-Q). *Can J Sport Sci* 17(4), 338-45 (1992).
- Craig C, Marshall AL, Sjoström M et al: International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 35(8), 1381-95 (2003).
- van Vugt JP, Sieling S, Piet KK et al.: Quantitative assessment of daytime motor activity provides a responsive measure of functional decline in patients with Huntington's disease. *Mov Disord* 16(3), 481-8 (2001).
- van Vugt JP, Piet KK, Vink LJ et al: Objective assessment of motor slowness in Huntington's disease: clinical correlates and 2-year follow-up. *Mov Disord* 19(3), 285-297 (2004).
- Busse ME, van Deursen RW, Wiles CM: Activity indices for measuring mobility in neurologically impaired patients. *Journal of neurology neurosurgery and psychiatry* 74(10), 1459-1459 (2003).
- Busse ME, Wiles CM, Rosser AE: Mobility and falls in people with Huntington's disease. *Journal of neurology neurosurgery and psychiatry* 80(1), 88-90 (2009).
- Noble BJ, Robertson RJ: Perceived Exertion. Champaign, IL: Human Kinetics (1996).
- Day ML, McGuigan MR, Brice G, Foster C: Monitoring exercise intensity during resistance training using the session RPE scale. *Journal of Strength and Conditioning Research*, 8(2), 53-58 (2004).
- Turner-Stokes L: Goal attainment scaling (GAS) in rehabilitation: a practical guide. *Clin Rehabil*, 23(4), 362-70 (2009).
- Enright PL: The six-minute walk test. *Respiratory Care* 48(8), 783-785 (2003).
- West C, Bowen A, Hesketh A, Vall A: Interventions for motor apraxia following stroke. *Cochrane Database Syst Rev*, 2008(1), CD004132 (2008).
- Donkervoort M, Dekker J, Stehmann-Saris FC: Efficacy of strategy training in left hemisphere stroke patients with apraxia: a randomized clinical trial. *Neuropsychological Rehabilitation*, 11(5), 549-566 (2011).
- Melton AK, Bourgeois MS: Training compensatory memory strategies via telephone for persons with TBI. *Aphasiology*, 2005, 19,353-364.
- Leng TR, Woodward MJ, Stokes MJ, Swan AV, Vaireing LA, Baker R: Effects of multisensory stimulation in people with Huntington's disease: a randomized controlled pilot study. *Clinical Rehabilitation*, 17(1), 30-41 (2003).
- Ho AK, Robbins AO, Walters SJ, Kaptoge S, Sahakian BJ, Barker RA: Health-related quality of life in Huntington's disease: a comparison of two generic instruments, SF-36 and SIP. *Mov Disord* 19(11), 1341-8 (2004).
- Watson MJ: Refining the ten-metre walking test for use with neurologically impaired people. *Physiotherapy* 88(7), 386-397 (2002).
- Podsiadlo D, Richardson S: The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*, 39(2), 142-148 (1991).
- Rao AK, Muratori L, Louis ED, Moskowitz CB, Marder KS: Clinical measurement of mobility and balance impairments in Huntington's disease: validity and responsiveness. *Gait Posture*, 29(3), 433-6 (2009).
- Whitney S, Marchetti GF, Morris LO, Sparto PJ: The reliability and validity of the four square step test for people with balance deficits secondary to vestibular disorder. *Archives of Physical Medicine and Rehabilitation*. 88, 99 (2008).
- VanHeugten CM, Dekker J, Deelman BG, Stehmann-Saris JC, Kinebanian A: A diagnostic test for apraxia in stroke patient: internal consistency and diagnostic value. *The Clinical Neuropsychologist* 13, 182 (1999).
- Whitney SL, Wrisley DM, Marchetti GF, Gee MA, et al. Clinical measurement of sit-to-stand performance in people with balance disorders: validity of data for the five-times-sit-to-stand test. *Phys Ther* 2005;85(10):1034-1045.
- Lasker AG, Zee DS: Ocular motor abnormalities in Huntington's disease. *Vision Res* 37(24), 3639-45 (1997).
- Lemiere J, Decruyenaere M, Evers-Kiebooms G, Vandenbusche E, Dom R: Cognitive changes in patients with Huntington's disease (HD) and asymptomatic carriers of the HD mutation—a longitudinal follow-up study. *Journal of Neurology*, 251(8), 935-42 (2004).
- Grimbergen Y, Knol MJ, Bloem BR, Kremer BP, Roos RA, Munneke M: Falls and gait disturbances in Huntington's disease. *Mov Disord* 23(7), 970-6 (2008).
- Aubeeluck A, Wilson E: Huntington's disease. Part 1: essential background and management. *British Journal of Nursing* 17(3), 146-151 (2008).
- Louis ED, Lee P, Quinn L, Marder K: Dystonia in Huntington's disease: prevalence and clinical characteristics. *Mov Disord* 4(1), 95-101 (1999).
- Penney JB, Vonsattel JP, MacDonald ME, Gusella JF, Myers RH: CAG repeat number governs the development rate of pathology in Huntington's disease. *Ann Neurol* 41(5), 689-92 (1997).
- Rosenblatt A, Abbott MH, Gourley LM et al: Predictors of neuropathological severity in 100 patients with Huntington's disease. *Annals of Neurology* 54(4),488-93 (2003).
- Vonsattel JP, Keller C, Cortes Ramirez EP: Huntington's disease - neuropathology. *Handbook Clin Neurol* 100, 83-100 (2011).
- Busse M, Hughes G, Wiles CM, Rosser AE: Use of hand-held dynamometry in the evaluation of lower limb muscle strength in people with Huntington's disease. *J Neurol* 255(10), 1534-40 (2008).
- Tian J, Herdman SJ, Zee DS, Folstein SE: Postural stability in patients with Huntington's disease. *Neurology* 42(6), 1232-8 (1992).
- Panzera R, Salomonczyk D, Progovosky E, et al: Postural deficits in Huntington's disease when performing motor skills involved in daily living. *Gait Posture* 33(3), 457-61 (2011).
- Rao A, Muratori L, Louis L, Moskowitz C, Marder K: Spectrum of gait impairments in presymptomatic and symptomatic Huntington's disease. *Mov Disord* 23(8), 1100-7 (2008).
- Hausdorff, JM: Gait variability and basal ganglia disorders: stride-to-stride variations of gait cycle timing in Parkinson's disease and Huntington's disease. *Mov Disord* 13(3), 428-437 (1998).
- Delval A, Krystkowiak P, Delliaux M et al: Effect of external cueing on gait in Huntington's disease. *Mov Disord* 23(10), 1446-52 (2008).
- Delval A, Krystkowiak P, Delliaux M et al: Role of attentional resources on gait performance in Huntington's disease. *Mov Disord* 23(5), 684-9 (2008).
- O'Donnell BF: Visual perception in prediagnostic and early stage Huntington's disease. *J Int Neuropsychol Soc* 14(3), 446-53 (2008).
- Hicks, SL, Robert MP, Golding CV, Tabrizi SJ, Kennard C: Oculomotor deficits indicate the progression of Huntington's disease. *Prog Brain Res* 171, 555-8 (2008).
- Peacock IW: A physical therapy program for Huntington's disease patients. *Clinical Management in Physical Therapy* 7(1), 22-23 (1987).
- Zinzi P, Salmasso D, De Grandis R et al: Effects of an intensive rehabilitation programme on patients with Huntington's disease: a pilot study. *Clin Rehabil* 21(7), 603-613 (2007).
- Thaut MH, Miltner R, Lange HW, Hurt CP, Hoemberg V: Velocity modulation and rhythmic synchronization of gait in Huntington's disease. *Mov Disord* 14(5), 808-819 (1999).
- Churchyard AJ, Morris ME, Georgiou N, Chiu E, Cooper R, Iansek R: Gait dysfunction in Huntington's disease: parkinsonism and a disorder of timing. Implications for movement rehabilitation. *Advances in Neurology* 87, 375-85 (2001).
- Kloos A, Kegelmeyer D, Kostyk S: The Effects of Assistive Devices on Gait Measures in Huntington's Disease. *Neurotherapeutics* 6(1), 209-210 (2009).
- Binley B, Morris ME, Denisenko S: Physiotherapy for people with movement disorders arising from basal ganglia dysfunction. *New Zealand Journal of Physiotherapy* 31(2), 94-100 (2003).
- Huntington Study Group. Unified Huntington's Disease Rating Scale: reliability and consistency. *Mov Disord* 1996; 11: 136-142.
- Byltsma FW, Rothind J, Hall MR, Folstein SE, Brandt J. Assessment of adaptive functioning in Huntington's disease. *Mov Disord* 1993;8 (2):183-190.
- Powell LE, Myers AM: The Activities-specific Balance Confidence (ABC) Scale. *J Gerontol A Biol Sci Med Sci* 50A(1), M28-M34 (1995).
- Rao A et al: Clinical measurement of mobility and balance impairments in Huntington's disease: validity and responsiveness. *Gait Posture* 29(3), 433, 6 (2009).
- Berg KO et al: Measuring balance in the elderly: validation of an instrument. *Can J Public Health* 83 Suppl 2, S7-11 (1992).
- Tinetti ME: Performance-oriented assessment of mobility problems in elderly patients. *J Am Geriatr Soc* 34(2), 119-126 (1986).
- Kloos AD et al: Fall risk assessment using the Tinetti mobility test in individuals with Huntington's disease. *Mov Disord* 25(16), 2838-44 2010.
- Rao AK, Quinn L, Marder KS: Reliability of spatiotemporal gait outcome measures in Huntington's disease *Mov Disord* 20(8), 1033-1037 (2005).
- Aziz NA : Weight loss in Huntington disease increases with higher CAG repeat number. *Neurology* 71(19), 1506-13 (2008).
- Wong D: Pain in children: comparison of assessment scales. *Pediatr Nurs* 14, 9-17 (1998).
- Jackson J: Specific treatment techniques. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds). Churchill Livingstone Elsevier, 243-266 (2011).
- Haas BA: Physical Activity and Exercise in Neurological Rehabilitation. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds). Churchill Livingstone Elsevier, 349-365 (2011).
- Kunkel, DSE: Falls and their management. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds). Churchill Livingstone Elsevier, 383-397 (2011).
- Jones F: Self management. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds), 367-387 (2011).
- Watson, P: Pain management in Neurological Rehabilitation. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds), 331-338 (2011).
- Folstein MF, Robins LN, Helzer JE: The Mini-Mental State Examination. *Arch Gen Psychiatry* 40(7), 812 (1983).
- Kilbride CC : Physical Management of Altered Tone and Movement. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds), 289-318 (2011).
- Bruton A: Respiratory Management in Neurological Rehabilitation. In *Physical Management for Neurological Disease*, M Stokes & E Stokes (Eds), 319-330 (2011).
- Zarit SH, Reever KE, Bach-Peterson J. Relatives of the impaired elderly: correlates of feelings of burden. *Gerontologist* 1980;20:649-55.
- Bott J: Guidelines for the physiotherapy management of the adult, medical, spontaneously breathing patient. *Thorax* 64(Suppl 1), i1-i52 (2009).
- Moxham J: Breathlessness, fatigue and the respiratory muscles. *Clinical Medicine, Journal of the Royal College of Physicians* 9(5), 448-52 (2009).
- Jones U, Enright S, Busse M: Management of respiratory problems in people with neurodegenerative conditions: A narrative review. *Physiotherapy* (in press).
- McCool FD, Rosen MJ: Nonpharmacologic airway clearance therapies: ACCP evidence-based clinical practice guidelines. *Chest* 129 (1 Suppl), 250S-259S (2006).
- Innocent DM, TF: Dysfunctional breathing, in *Physiotherapy for respiratory and cardiac problems*, PA Prasad (Ed), Churchill Livingstone: Oxford (2008).
- Yorkston KM, Millar RM, Klasner ER: Huntington's Disease. In *Management of Speech and Swallowing in Degenerative Disease*. KM, Yorkston KM, RM Millar & Stroud EA (Eds). PRO-ED: Austin TX, 139-154 (2004).
- Braden BJ Bergstrom N 1994 Predictive validity of the Braden scale for pressure sore risk in a nursing home. *Res Nurs Health* 17: 459-70
- European Pressure Ulcer Advisory panel (EPUAP and national Pressure Ulcer Advisory Panel (NPUAP). *Pressure Ulcer Treatment: Quick Reference Guide*. www.npuap.org

Huntington's disease

European Huntington's Disease Network Physiotherapy Working Group*

Please send your comments, suggestions and overall feedback to **Lori Quinn** QuinnL1@cardiff.ac.uk

Imprint :

© 2013 European Huntington's Disease Network,

**Chairman Prof. G.B. Landwehrmeyer,
Oberer Eselsberg 45/1, 89081 Ulm, Germany,**

www.euro-hd.net

Acknowledgements :

*Edited by Lori Quinn, EdD, PT and Monica Busse, PhD MSc (Med) BSc (Med) Hons BSc (Physio)
Initial versions of this document were developed by a sub-group physiotherapists: Lori Quinn, Monica Busse, Hanan Khalil, Una Jones, Angela Hall, Sue Armstrong.
Subsequent development and field testing completed by members of the EHDN Physiotherapy Working Group, with specific contributions by: Karin Bunnig, Maggie Broad, Katy DeBono, Camilla Ekwall, Hanne Fossmo, Nora Fritz, Karen Jones, Una Jones, Deb Kegelmeyer, Hanan Khalil, Anne Kloos , Rodolfo Vera, Alexandra Schuler, and Jessie van der Bent

The information contained in this brochure is subject to the European HD Network Liability Disclaimer which can be found at <http://www.euro-hd.net/html/disclaimer>.

– Please consult a doctor for medical advice.

– Except as otherwise noted this work is licensed under the **Creative Commons Attribution-No Derivative Works 3.0 Unported License**.